Child's Name  Avoid Triggers: (Check all that  Illness	☐ Cigarette/other smoke	□ Food:
☐ Emotions	☐ Exercise	☐ Allergies:
☐ Weather Changes	☐ Chemical odors	☐ Other:
Green Zone: Child breathing at best Well	Yellow Zone: Child not breathing at best Sick	Red Zone:  Danger Zone  Emergency
<ul> <li>sleeps through the night without coughing or wheezing</li> <li>has no early warning signs of an asthma flare-up</li> <li>plays actively</li> </ul>	<ul> <li>coughing or wheezing at night or at child care</li> <li>has early warning signs of a flare-up:</li> </ul>	<ul> <li>breathing is hard and fast</li> <li>coughing, short of breath, wheezing</li> <li>neck and chest "suck in" skin between ribs, above the breastbone and collarbone where</li> </ul>
Take Long-Term Control medications:  •	<ul> <li>has trouble doing usual activities/play,</li> <li>may self limit activities/ squat/hunch over</li> <li>decrease in appetite/difficulty drinking or taking a bottle.</li> </ul>	<ul> <li>breathing</li> <li>has trouble walking or talking</li> <li>stops activities</li> <li>unable to drink or take bottle</li> </ul>
Take quick-relief medicines 15 minutes before active playtime.	Take quick–relief medicines:  Adjust Long-Term Control medicines as follows until back in Green Zone:	Emergency Medicine Plan:  Call 911 if  no improvement 15 minutes after quick relief medication given and  nails or lips are blue  is having trouble walking or
Parent:	Activity Restrictions: Ozone Restrictions:	talking • cannot stop coughing
Physician:  Telephone:  Adapted by the NC Child Care Health Consultants Association	Call child's parent if:  child's symptoms do not improve or worsen 15 to 20 minutes after treatment  Call the physician if: parent not available	Physician Signature  Date: